

# The moduli space of local homogeneous 3-geometries†

Martin Rainer ‡

Gravitationsprojekt / Kosmologie, Institut für Mathematik,  
Universität Potsdam, PF 601553, D-14415 Potsdam, Germany  
e-mail: mrainer@aip.de

**Abstract:** For a canonical formulation of quantum gravity, the superspace of all possible 3-geometries on a Cauchy hypersurface of a 3 + 1-dimensional Lorentzian manifold plays a key role. While in the analogous 2 + 1-dimensional case the superspace of all Riemannian 2-geometries is well known, the structure of the superspace of all Riemannian 3-geometries has not yet been resolved at present.

In this paper, an important subspace of the latter is disentangled: The superspace of local homogenous Riemannian 3-geometries. It is finite dimensional and can be factored by conformal scale dilations, with the flat space as the center of projection. The corresponding moduli space can be represented by homothetically normalized 3-geometries.

By construction, this moduli space of the local homogenous 3-geometries is an algebraic variety. An explicit parametrization is given by characteristic scalar invariants of the Riemannian 3-geometry.

Although the moduli space is not locally Euclidean, it is a Hausdorff space. Nevertheless, its topology is compatible with the non-Hausdorffian topology of the space of all Bianchi-Lie algebras, which characterize the moduli modulo differences in their anisotropy.

† This lecture is financially supported by the DAAD (Germany) and the IPM (Iran).

‡Present address: Institute for Studies in Physics and Mathematics  
Farmanieh Bld., Tehran P.O.Box 19395-5531, Tehran, Iran  
e-mail: mrainer@physics.ipm.ac.ir

## 1. Introduction

The canonical formalism of (quantum) gravity uses a topological  $d + 1$  decomposition of the  $(d + 1)$ -dimensional Lorentzian manifold  $M^{d+1} = M^1 \times M^d$ , into a time manifold  $M^1$  and a smooth topological  $d$ -space  $M^d$ , which for any  $t \in M^1$  is a Riemannian manifold  $M^d(t)$ . The action for pure Einstein gravity on  $M^{d+1}$  then becomes

$$\begin{aligned} S &= \int_{M^{d+1}} \sqrt{|\det^{(d+1)}g|} \, {}^{(d+1)}R \, d^{d+1}x \\ &= \int_{M^1} \int_{M^d(t)} N \sqrt{\det g(t)} \, \mathcal{L}(t) \, d^d x \, dt, \end{aligned} \quad (1.1)$$

which is an integral over all Riemannian hypersurfaces  $M^d(t)$ , where  $\mathcal{L} := R + \text{tr}(K^2) - (\text{tr}K)^2$ . On each  $M^d(t)$  the Riemannian metric  $g(t)$  yields intrinsic Ricci scalar curvature  $R(t)$  and the extrinsic curvature form  $K(t)$ . The latter is the second fundamental form with components

$$K_{\alpha\beta} = -\frac{1}{2N} \left( \frac{\partial g_{\alpha\beta}}{\partial t} - N_{\alpha;\beta} - N_{\beta;\alpha} \right), \quad \alpha, \beta = 1, \dots, d, \quad (1.2)$$

where  $N$  and  $N_\alpha$  are the lapse and shift functions, respectively. Note that, throughout this paper, any Riemannian metric  $g$  is per definition sign normalized to  $\det g > 0$ . Using the conjugate momenta

$$\pi = \frac{1}{\sqrt{\det g}} \frac{\delta S}{\delta \dot{g}}, \quad (1.3)$$

the Hamiltonian density becomes

$$\begin{aligned} \mathcal{H} &= \pi^{\alpha\beta} g_{\alpha\beta} - \mathcal{L} \\ &= N \left[ \text{tr}(\pi^2) - \frac{1}{d-1} (\text{tr}\pi)^2 - R \right] + 2N_\alpha \pi^{\alpha\beta}{}_{;\beta} \\ &= N \mathcal{C} + 2N_\alpha \mathcal{C}^\alpha, \end{aligned} \quad (1.4)$$

where  $\mathcal{C}$  and  $\mathcal{C}^\alpha$  are the super-Hamiltonian and super-momentum constraints, enforcing reparametrization invariance of  $M^1$  and  $M^d(t)$ , respectively. In particular, the possible reparametrizations of any  $M^d$  are given by the connected component  $\text{Diff}_0(M^d)$  of its diffeomorphism group  $\text{Diff}(M^d)$ .

Let  $\text{Met}(M^d)$  be the space of all Riemannian metrics on  $M^d$ . Then the Wheeler superspace of  $d$ -geometries is given as

$$\text{Geom}(M^d) := \frac{\text{Met}(M^d)}{\text{Diff}(M^d)}. \quad (1.5)$$

In the following,  $g \in \text{Geom}(M^d)$  is always understood to denote the geometry as diffeomorphism invariant class of the corresponding metrics. In generic regions,  $\text{Geom}(M^d)$  can be equipped with an ultralocal metric, given as a reparametrization invariant form with components

$$G_{\alpha\beta\mu\nu}(x, x') := \frac{\delta^d(x - x')}{\sqrt{\det g}} (a g_{\alpha\mu} g_{\beta\nu} + b g_{\alpha\nu} g_{\beta\mu} - c g_{\alpha\beta} g_{\mu\nu}), \quad (1.6)$$

$\alpha, \beta, \mu, \nu = 1, \dots, d$ . For coefficients  $(a, b, c) = (1, 0, \frac{1}{d-1})$  this is the DeWitt metric, used e.g. in [1]. A different choice, namely  $(a, b, c) = (1, 1, 1)$ , was taken in [2, 3, 4]. However, we should not expect such a metric (1.6) to exist on all of  $\text{Geom}(M^d)$ , for the following reason: It is known that the action of  $\text{Diff}(M^d)$  on  $\text{Met}(M^d)$  is not free in general. So with the natural quotient topology,  $\text{Geom}(M^d)$  is not a manifold. At certain symmetrical metrics the quotient has singularities. However, with some techniques from algebraic geometry (in [5] also applied to ADE singularities), a first non-singular resolution of  $\text{Geom}(M^d)$  was found in [6], and the minimal non-singular resolution  $\text{Geom}_o(M^d)$  was finally constructed in [7].

Although the classical action (1.1) is given as the integral over a path of  $d$ -geometries  $g \in \text{Geom}(M^d)$ , for its quantization the superspace  $\text{Geom}(M^d)$  is not the appropriate configuration space, because the Euclidean path integral would then contain a divergent integration over the conformal modes of the geometry. In [8] it was suggested to use a Lorentzian path integral instead. However, the superspace  $\text{Geom}(M^d)$  could also be regularized, if one succeeds to factor out the conformal modes. Indeed, for  $d = 2$  this has been done successfully in [1]. The superspace  $\text{Geom}(M^2)$  of all Riemannian 2-geometries  $g$  on  $M^2$  can be disentangled, into the conformal deformations  $\text{Conf}(M^2)$  of any Riemannian geometry  $g$ , and the moduli space  $\mathcal{M}(M^2)$  of all conformal equivalence classes  $[g]$ :

$$\text{Geom}(M^2) = \text{Conf}(M^2) \times \mathcal{M}(M^2). \quad (1.7)$$

Since any Riemannian 2-manifold  $M^2$  is conformally flat, all local data on  $M^2$  is actually contained in  $\text{Conf}(M^2)$ . However, for  $d > 2$ , the local data of the manifold  $M^d$  is partially also contained in the moduli space<sup>1</sup>

$$\mathcal{M}(M^d) := \frac{\text{Geom}(M^d)}{\text{Conf}(M^d)} \quad (1.8)$$

<sup>1</sup>Note that mathematically this term applies well to the present case, although some physicists use it with a more limited understanding.

of Riemannian  $d$ -geometries  $\text{Geom}(M^d)$  modulo conformal transformations  $\text{Conf}(M^d)$ . Therefore, for  $d > 2$ , it makes sense to separate the local data in  $\mathcal{M}(M^d)$ , neglecting all global properties of  $M^d$ . When  $M^d$  is considered only locally, the mapping class group  $\text{Diff}(M^d)/\text{Diff}_0(M^d)$  is negligible, and  $\text{Geom}(M^d)$  is really the classical configuration space. For a *local homogeneous* manifold  $M^d$ , all geometrical data is given localized at just one arbitrarily chosen point  $x$  of  $M^d$ , on a neighbourhood  $U_\epsilon(x)$ , with infinitesimal limit  $\epsilon \rightarrow 0$ . Therefore we denote *local homogeneous Riemannian  $d$ -geometries* just as  $\text{Geom}(d)$ . *Homogeneous conformal transformations*, denoted as  $\text{Conf}(d)$ , leave the unique flat Riemannian geometry  $I^d$  invariant. Therefore, we consider in the following the *moduli space of local homogeneous  $d$ -geometries* as defined by

$$\mathcal{M}(d) := \frac{\text{Geom}(d)}{\text{Conf}(d)} \setminus \{I^d\}. \quad (1.9)$$

While for dimension  $d \geq 4$  the moduli space  $\mathcal{M}(d)$  seems still to be exceedingly complicated, the structure of the moduli space  $\mathcal{M}(3)$  of local homogeneous Riemannian 3-geometries is resolved in the present paper.

Sec. 2 resumes the local properties of (Riemannian) homogeneous 3-geometries. In Sec. 3 the moduli space  $\mathcal{M}(3)$  is constructed as an algebraic variety, parametrized by scalar geometric invariants. Sec. 4 then shows the consistency of the locally non-Euclidean Hausdorff topology of  $\mathcal{M}(3)$  with a topological Morse-like potential on the space  $K^3$  of Bianchi Lie isometries of the moduli. Finally, Sec. 5 gives a short discussion of the results.

## 2. Local properties of homogenous 3-geometries

Per definition, a homogeneous geometry admits a transitive action of its isometry group. The Kantowski-Sachs (KS) spaces are the only homogeneous Riemannian 3-spaces not admitting a simply transitive subgroup of their isometry group. The Lie algebra of the latter is  $\mathbb{R} \oplus \text{IX}$ , the Abelian extension of the Bianchi Lie algebra  $\text{IX} = \text{so}(3)$ . Any KS space can be obtained as a specific limit of Bianchi IX spaces. (Globally, a hyper-cigar like 3-ellipsoid of topology  $S^3$  is stretched to infinite length, becoming a hyper-cylinder  $S^2 \times \mathbb{R}$ .) All other homogeneous Riemannian 3-spaces have a simply transitive isometry subgroup, corresponding to one of the Bianchi Lie algebras. Therefore, the following considerations can be restrict to homogenous 3-geometries of Bianchi type, nevertheless yielding finally a classification of *all* local homogeneous Riemannian 3-geometries. Let the *characteristic isometry* of a local homogeneous 3-geometry be defined to be IX for KS spaces or its Bianchi type otherwise.

In the case of a transitive Lie isometry of Bianchi type, the Riemannian geometry can be soldered to an orthogonal frame spanned by the Lie algebra generators  $e_i$  in the tangent space, i.e.

$$g_{\mu\nu} = e_\mu^a e_\nu^b g_{ab}, \quad \mu, \nu = 1, 2, 3, \quad (2.1)$$

where  $e^a = e_\mu^a dx^\mu = g^{ai} e_i$ ,  $e_i = e_i^\mu \frac{\partial}{\partial x^\mu}$ , and  $g^{ab} g_{ij} = \delta_i^a \delta_j^b$ , with the constant metric

$$(g_{ab}) = \begin{bmatrix} e^s & 0 & 0 \\ 0 & e^{s+w-t} & 0 \\ 0 & 0 & e^{s-t} \end{bmatrix}. \quad (2.2)$$

Here  $s$  fixes the overall scale, while  $t$  and  $w$  parametrize the anisotropies related respectively to the  $e_1$  and  $e_2$  direction (preserving isotropy in the respective orthogonal planes).

The data which characterizes the *local* geometrical structure of a homogeneous Riemannian manifold  $M^3$  of characteristic Bianchi isometry can be rendered in form of (i) the *local scales* of (2.2), (ii) the *covariant derivatives*

$$De^k = e_{i;j}^k e^i e^j := e_{\alpha;\beta}^k dx^\alpha dx^\beta \quad (2.3)$$

of the dual generators  $e^k$  in the cotangent frame, and iii) the corresponding *Bianchi Lie algebra*, represented as

$$[e_i, e_j] = C_{ij}^k e_k. \quad (2.4)$$

The bracket  $[\cdot, \cdot]$  defines a Lie algebra, iff the structure constants satisfy the antisymmetry condition

$$C_{[ij]}^l = 0, \quad (2.5)$$

and the Jacobi condition

$$C_{[ij}^l C_{kl]}^m = 0, \quad (2.6)$$

with nondegenerate antisymmetric indices  $i, j, k = 1, 2, 3$ . With (2.5) a Lie algebra is already completely described by the  $2 \times 2$ -matrices  $C_{<i>}$ ,  $i = 1, 2, 3$ , each with components  $C_{ij}^k$ ,  $j, k = 1, 2$ . But this description may still carry redundancies: The space of all sets  $\{C_{ij}^k\}$  satisfying the Lie algebra conditions (2.5) and (2.6) is a subvariety  $W^3 \subset \mathbb{R}^9$ .  $\text{GL}(3)$  basis transformations act on a given set of structure constants as tensor transformations:

$$C_{ij}^k \rightarrow \tilde{C}_{ij}^k := (A^{-1})_h^k C_{fg}^h A_i^f A_j^g \quad \forall A \in \text{GL}(3). \quad (2.7)$$

On  $W^3$  this yields a natural equivalence relation  $C \sim \tilde{C}$ , defined by

$$C_{ij}^k \sim \tilde{C}_{ij}^k : \Leftrightarrow \exists A \in \text{GL}(3) : \tilde{C}_{ij}^k = (A^{-1})_h^k C_{fg}^h A_i^f A_j^g. \quad (2.8)$$

The associated projection

$$\pi : \begin{cases} W^3 & \rightarrow K^3 := W^3/\text{GL}(3) \\ C & \mapsto [C] \end{cases} \quad (2.9)$$

yields just the space  $K^3$  of all Bianchi Lie algebras as quotient space.

The present choice of the real parameters  $s, t, w$  for datum (i) simplifies the following calculations in a specific triad basis for data (ii) and (iii), chosen in consistency with the representations of [9, 10]. The basis is given by matrices  $(e_\alpha^a)$ , with anholonomic rows  $a = 1, 2, 3$  indexing the generators of the algebra, and holonomic coordinate columns  $\alpha = 1, 2, 3$ . The coordinates will also be denoted as  $x := x^1$ ,  $y := x^2$ ,  $z := x^3$ . For the different Bianchi types the matrices  $(e_\alpha^a)$  take the following form:

Bianchi I:

$$(e_\alpha^a) = \begin{bmatrix} 1 & & \\ & 1 & \\ & & 1 \end{bmatrix}, \quad (2.10)$$

Bianchi II:

$$(e_\alpha^a) = \begin{bmatrix} 1 & -z & \\ 0 & 1 & \\ & & 1 \end{bmatrix}, \quad (2.11)$$

Bianchi IV:

$$(e_\alpha^a) = \begin{bmatrix} 1 & & \\ & e^x & 0 \\ & xe^x & e^x \end{bmatrix}, \quad (2.12)$$

Bianchi V:

$$(e_\alpha^a) = \begin{bmatrix} 1 & & \\ & e^x & \\ & & e^x \end{bmatrix}, \quad (2.13)$$

Bianchi VI<sub>h</sub>,  $h = A^2$ :

$$(e_\alpha^a) = \begin{bmatrix} 1 & & \\ & e^{Ax} \cosh x & -e^{Ax} \sinh x \\ & -e^{Ax} \sinh x & e^{Ax} \cosh x \end{bmatrix}, \quad (2.14)$$

Bianchi VII<sub>h</sub>,  $h = A^2$ :

$$(e_\alpha^a) = \begin{bmatrix} 1 & & \\ & e^{Ax} \cos x & -e^{Ax} \sin x \\ & e^{Ax} \sin x & e^{Ax} \cos x \end{bmatrix}, \quad (2.15)$$

Bianchi VIII:

$$(e_\alpha^a) = \begin{bmatrix} \cosh y \cos z & -\sin z & 0 \\ \cosh y \sin z & \cos z & 0 \\ \sinh y & 0 & 1 \end{bmatrix}, \quad (2.16)$$

Bianchi IX:

$$(e_\alpha^a) = \begin{bmatrix} \cos y \cos z & -\sin z & 0 \\ \cos y \sin z & \cos z & 0 \\ -\sin y & 0 & 1 \end{bmatrix}. \quad (2.17)$$

For each of the lines  $\{\text{VI}_h\}$  and  $\{\text{VII}_h\}$  the parameter range is given by  $\sqrt{h} = A \in [0, \infty[$ . Let us also remind that,  $\text{III} := \text{VII}_1$ .

The structure constants can be reobtained from  $ds^2$  and the triad by

$$C_{ijk} = ds^2([e_i, e_j], e_k), \quad C_{ij}^k = C_{ijr} g^{rk}. \quad (2.18)$$

The metrical connection coefficients are determined as

$$\Gamma_{ij}^k = \frac{1}{2} g^{kr} (C_{ijr} + C_{jri} + C_{irj}). \quad (2.19)$$

W.r.t. the triad basis, the curvature operator is defined as

$$\mathfrak{R}_{ij} := \nabla_{[e_i, e_j]} - (\nabla_{e_i} \nabla_{e_j} - \nabla_{e_j} \nabla_{e_i}), \quad (2.20)$$

the Riemann tensor components are

$$R_{hijk} := \langle e_h, \mathfrak{R}_{ij} e_k \rangle, \quad (2.21)$$

and the Ricci tensor is

$$R_{ij} := R_{ikj}^k = \Gamma_{ij}^f \Gamma_{fe}^e - \Gamma_{ie}^f \Gamma_{fj}^e + \Gamma_{if}^e \Gamma_{ej}^f. \quad (2.22)$$

From (2.22) we may form the following scalar invariants of the geometry: The Ricci curvature scalar

$$R := R^i_i, \quad (2.23)$$

the sum of the squared eigenvalues

$$N := R^i_j R^j_i, \quad (2.24)$$

the trace-free scalar

$$S := S^i_j S^j_k S^k_i = R^i_j R^j_k R^k_i - RN + \frac{2}{9} R^3, \quad (2.25)$$

where  $S^i_j := R^i_j - \frac{1}{3} \delta^i_j R$ , and, related to the York tensor,

$$Y := R_{ik;j} g^{il} g^{jm} g^{kn} R_{lm;n}, \quad \text{with} \quad (2.26)$$

$$\begin{aligned} R_{ij;k} &:= e_i^\alpha e_j^\beta e_k^\gamma R_{\alpha\beta;\gamma} \\ &= e_{\alpha;\beta}^l e_m^\alpha e_k^\beta (\delta_i^m \delta_j^n + \delta_i^n \delta_j^m) R_{lmn}. \end{aligned}$$

The 4 scalar invariants above characterize a local homogeneous Riemannian 3-geometry. It satisfies  $N = 0$ , iff it is the unique flat  $\text{I}^3$ . Besides the transitive Bianchi I isometry, it also admits a left-invariant (but not transitive) Bianchi  $\text{VII}_0$  action on its 2-dimensional hyperplanes (cf. [11, 12]).

### 3. Explicit construction of the moduli space $\mathcal{M}(3)$

Let  $\text{I}^3$  be a center of projection for all other geometries of  $\text{Geom}(3)$ . All these non-flat Bianchi or KS types satisfy  $N \neq 0$ . The invariant  $N$  then parametrizes (like  $e^{-2s}$ ) a 3-geometry's homogeneous conformal scale in  $\text{Conf}(3)$ . A homogeneous conformal transformation is just given as homothetic rescaling

$$g_{ij} \rightarrow \sqrt{N} g_{ij}, \quad (3.1)$$

yielding the following normalized invariants, which depend only on the homogeneously conformal class of the geometry:

$$\hat{N} := 1, \quad \hat{R} := R/\sqrt{N}, \quad \hat{S} := S/N^{3/2}, \quad \hat{Y} := Y/N^{3/2}. \quad (3.2)$$

For a non-flat Riemannian 3-space, the invariant  $\hat{Y}$  vanishes, iff the 3-geometry is conformally flat. A general conformal transformation is not necessarily homogeneous. Hence there may exist homogeneous spaces, which are in the same conformal class, but in different homogeneously conformal classes. Note that a rescaling (3.1) does not change the Bianchi or KS type of isometry.

The moduli space  $\mathcal{M}(3)$  is given by classifying space  $\text{Geom}(3)$  of non-flat local homogeneous Riemannian 3-geometries modulo homogeneous conformal transformations  $\text{Conf}(3)$  (3.1). Since the flat Bianchi I geometry, as  $\text{Conf}(3)$ -invariant center of the projection, has been excluded, the moduli space  $\mathcal{M}(3)$  can be parametrized by the invariants  $\hat{R}$ ,  $\hat{S}$  and  $\hat{Y}$ , given for each fixed Bianchi type as a function of the anisotropy parameters  $t$  and  $w$ . For the non-flat Bianchi Lie algebras these invariants are listed explicitly in [13].

$\mathcal{M}(3)$  can also be embedded into a minimal cube, spanned by  $\hat{R}/\sqrt{3}$ ,  $\sqrt{6}\hat{S} \in [-1, 1]$  and  $2 \tanh \hat{Y} \in [0, 2]$ . Fig. 1 describes points of the moduli space which are of Bianchi type VI and VII or lower. Fig. 2 shows likewise points for Bianchi type VIII and IX.

A projection of  $\mathcal{M}(3)$  to the  $\hat{R}$ - $\hat{S}$ -plane was already considered in [14]. For a homogeneous space with 2 equal Ricci eigenvalues the corresponding point in the  $\hat{R}$ - $\hat{S}$ -plane lies on a double line  $L_2$ , which has a range defined by  $|\hat{R}| \leq \sqrt{3}$  and satisfies the algebraic equation

$$162\hat{S}^2 = (3 - \hat{R}^2)^3. \quad (3.3)$$

All other algebraically possible points of the  $\hat{R}$ - $\hat{S}$ -plane lie inside the region surrounded by the line  $L_2$ . At the branch points  $\hat{R} = \pm\sqrt{3}$  of  $L_2$  all Ricci eigenvalues are equal. These homogeneous spaces possess a 6-dimensional isometry group. Homogeneous spaces possessing a 4-dimensional isometry group are represented by points on  $L_2$ . If one Ricci eigenvalue equals  $R$ , i.e. if there exists a pair  $(a, -a)$  of Ricci eigenvalues, the corresponding point in the  $\hat{R}$ - $\hat{S}$ -plane lies on a line  $L_{+-}$ , defined by the range  $|\hat{R}| \leq 1$  and the algebraic equation

$$\hat{S} = \frac{11}{9} \hat{R}^3 - \hat{R}. \quad (3.4)$$

In the case that one eigenvalue of the Ricci tensor is zero, the corresponding point in the  $\hat{R}$ - $\hat{S}$ -plane lies on a line  $L_0$ , defined by the range  $|\hat{R}| \leq \sqrt{2}$  and the algebraic equation

$$\hat{S} = \frac{\hat{R}}{2} (1 - \frac{5}{9} \hat{R}^2). \quad (3.5)$$

At the branch points of the curve  $L_2$  the Ricci tensor has a triple eigenvalue, which is negative for geometries of Bianchi type V, and positive for type IX geometries with parameters  $(t, w) = (0, 0)$ . These constant curvature geometries are all conformally flat with  $\hat{Y} = 0$ . Besides the flat Bianchi I geometry, the remaining conformally flat spaces with  $\hat{Y}$  are the KS space  $(\hat{R}, \hat{S}, \hat{Y}) = (\sqrt{2}, -\frac{\sqrt{2}}{18}, 0)$  and, point reflected, the Bianchi type III<sub>c</sub>, corresponding to the initial point of a Bianchi III line segment ending at the Bianchi II point in Fig. 1.

The point  $(-1, 0, 0)$  of Fig. 1 admits both types, Bianchi V and VII<sub>h</sub> with  $h > 0$ . Nevertheless, this point corresponds only to one homogeneous space, namely the space of constant negative curvature. This is possible, because this space has a 6-dimensional Lie group, which contains the Bianchi V and VII<sub>h</sub> subgroups. Note that in the flat limit  $V \rightarrow I$ , the additional Bianchi groups VII<sub>h</sub> change with  $h \rightarrow 0$ .

Similarly, the Bianchi III points of Fig. 1 lie on the curve  $L_2$ . However, these points are also of Bianchi type VIII. In fact, each of them correspond to one homogeneous geometry only. However, the latter admits a 4-dimensional isometry group, which has two 3-dimensional subgroups, namely Bianchi III and VIII, both containing the same 2-dimensional non-Abelian subgroup.

Altogether, the moduli space  $\mathcal{M}(3)$  of local homogeneous Riemannian 3-spaces is a  $T_2$  (Hausdorff) space. But it is not a topological manifold: The line of VII<sub>0</sub> moduli is a common boundary of 3 different 2-faces, namely that of the IX moduli, that of the VIII moduli, and with  $h \rightarrow 0$  that of all moduli of type VII<sub>h</sub> with  $h > 0$ . With  $\mathcal{M}(3)$  also  $\text{Geom}(3)$  is not locally Euclidean; rather both are stratifiable varieties.  $\text{Geom}(3)$  is a projective cone with sections  $\mathcal{M}(3)$  and the flat geometry  $I^3$  as singularity. In the following Section it is shown, that the topology of  $\mathcal{M}(3)$  is consistent with the natural topology on the space  $K^3$  of Bianchi Lie algebras, which may provide a Morse-like isometry potential on  $\mathcal{M}(3)$ .

#### 4. A potential on the isometry components of $\mathcal{M}(3)$

Let us now examine the 3-dimensional characteristic isometries of the moduli in more detail. The space  $K^3$  of Bianchi Lie algebras is naturally equipped with the quotient topology  $\kappa^3$ , generated by the projection  $\pi$  from the subspace topology on  $W^3 \subset \mathbb{R}^9$ . Below it will become clear that  $\kappa^3$  does not satisfy the separation axiom  $T_1$ , i.e.  $K^3$  contains non-closed points, or in other words, there is a non-trivial transition from  $A$  to some  $B \neq A$  in the closure  $\text{cl}\{A\}$  of  $A$ . Non-trivial ( $A \neq B$ ) transitions are special limits, which exist only due to the non- $T_1$  property of  $\kappa^3$ . This property implies that  $\kappa^3$  is not Hausdorffian (i.e. the separation axiom  $T_2$  fails with  $T_1$ ). Here a *transitions* from  $A$  to  $B$  is defined by

$$A \geq B : \Leftrightarrow B \in \text{cl}\{A\}. \quad (4.1)$$

By this definition, transitions are transitive and yield a natural partial order. A transition  $A \geq B$  is non-trivial, iff  $A > B$ .

In order to find the topology  $\kappa^3$  and all its transitions in  $K^3$ , an explicit description of the Bianchi Lie algebras is useful. In fact, it can be given in terms of the nonvanishing matrices  $C_{<i>}$ ,  $i = 1, 2, 3$ . Furthermore, this representation can be normalized modulo an overall scale of the basis  $e_1, e_2, e_3$ , and  $C_3$  can be chosen in some normal form ([15, 16] use the Jordan normal form).

In the *semisimple* representation category, there are only the simple Lie algebras VIII  $\equiv \text{so}(1, 2) = \text{su}(1, 1)$  and IX  $\equiv \text{so}(3) = \text{su}(2)$ , given by

$$\begin{aligned} C_{<3>}(\text{VIII}) &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ C_{<1>}(\text{VIII}) &= -C_{<2>}(\text{VIII}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ C_{<3>}(\text{IX}) &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ C_{<1>}(\text{IX}) &= -C_{<2>}(\text{IX}) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \end{aligned} \quad (4.2)$$

All other algebras are in the *solvable* representation category. They all have an Abelian ideal  $\text{span}\{e_1, e_2\}$ . Hence, with vanishing  $C_{<1>} = C_{<2>} = 0$ , they are described by  $C_{<3>}$  only. The "inhomogeneous" algebras VI<sub>0</sub>  $\equiv \text{e}(1, 1) = \text{iso}(1, 1)$  (local isometry of a Minkowski plane) and VII<sub>0</sub>  $\equiv \text{e}(2) = \text{iso}(2)$  (local isometry of an Euclidean plane) are determined by

$$C_{<3>}(\text{VI}_0) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad C_{<3>}(\text{VII}_0) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (4.3)$$

It holds  $C_{<3>}(\text{VIII}) = C_{<3>}(\text{VII}_0) = C_{<3>}(\text{IX})$ . Furthermore,  $C_{<1>}(\text{VIII}) = C_{<3>}(\text{VI}_0)$  and  $C_{<1>}(\text{IX}) = C_{<3>}(\text{VII}_0)$ . So we find both transitions VIII  $\leq$  VII<sub>0</sub> and IX  $\leq$  VII<sub>0</sub>, but only VIII  $\leq$  VI<sub>0</sub>. These inhomogeneous algebras are endpoints  $h = 0$  of two 1-parameter sets of algebras, VI<sub>h</sub> and VII<sub>h</sub>, for  $h > 0$  given respectively as

$$C_{<3>}(\text{VI}_h) = \begin{pmatrix} h & 1 \\ 1 & h \end{pmatrix}, \quad C_{<3>}(\text{VII}_h) = \begin{pmatrix} h & 1 \\ -1 & h \end{pmatrix}. \quad (4.4)$$

Note also that the 1-parameter set of algebras VI<sub>h</sub>,  $0 \leq h < \infty$  contains an exceptional decomposable point III := VI<sub>1</sub>.  $C_{<3>}(\text{III})$  has exactly 1 zero eigenvalue, while for all other algebras VI<sub>h</sub> and VII<sub>h</sub>,  $0 \leq h < \infty$  the matrix  $C_{<3>}$  has two different non-zero eigenvalues, which become equal only in the limit  $h \rightarrow \infty$ . If the geometric multiplicity of this limit is 1, then the latter corresponds to the Bianchi Lie algebra IV, representable with

$$C_{<3>}(\text{IV}) = \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}. \quad (4.5)$$

By *geometric specialization* of the algebra IV the geometric multiplicity of its eigenvalue is increases to 2, yielding the pure vector type algebra V, given by

$$C_{<3>}(\text{V}) = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}. \quad (4.6)$$

By *algebraic specialization* of the algebra IV the eigenvalue becomes zero, yielding the Heisenberg algebra II, given by

$$C_{<3>}(\text{II}) = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}. \quad (4.7)$$

This algebra can also be reached by via a direct transition from any of the algebras VI<sub>h</sub> and VII<sub>h</sub>,  $0 \leq h < \infty$ . Finally both, geometric specialization of II and algebraic specialization of V, yield the unique Abelian Lie algebra I, satisfying

$$C_{<i>}(\text{I}) = 0, \quad i = 1, 2, 3. \quad (4.8)$$

Altogether we have found all non-trivial transitions of  $\kappa^3$ . So the topological space  $(K^3, \kappa^3)$  may be described explicitly by a minimal graph, constructed like following: Let us associate an *arrow*  $A \rightarrow B$  to a pair of algebras  $A, B \in K^3$ , with  $A > B$ , such that there exists no  $C \in K^3$  with  $A > C > B$ . We call  $A$  the *source* and  $B$  the *target* of the arrow  $A \rightarrow B$ . Now we define a discrete *index* function  $J : K^3 \rightarrow \mathbb{N}_0$  as following: We start with the unique element I, to which we assign the minimal index  $J(\text{I}) = 0$ . Then, for  $i \in \mathbb{N}_0$ , we assign the index  $J(S) = i + 1$  to the source algebra  $S$  of any arrow pointing towards a target algebra  $T$  of index  $J(T) = i$ , until, eventually for some index  $J = i_{\max}$  there is no arrow to any target algebra  $T$  with  $J(T) = i_{\max}$ . Let us denote the subsets of all elements with index  $i$  as *levels*  $L(i) \subset K^3$ .

$K^3$  is directed towards its *minimal element*, the Abelian Lie algebra I, constituting the only *closed point*. Due to some points which are neither open nor closed,  $K^3$  is connected in the topology  $\kappa^3$ . *Open points* correspond to *locally rigid* Lie algebras  $C$ , i.e. those which cannot be deformed to some  $A \geq C$  with index  $J(A) > J(C)$ . In this sense, the open points in  $K^n$  are its *locally maximal elements*. Furthermore, (Hausdorff) *isolated open points*

correspond to *rigid* Lie algebras  $C$ , i.e. those which cannot be deformed to any  $A \in K^3$  with  $A \not\leq C$  and index  $J(A) \geq J(C)$ . In this sense, the isolated open points are the *isolated locally maximal elements*.

Fig. 3 shows on each horizontal level  $L(i)$  the algebras of equal index  $i$ , which are the sources for the level  $L(i-1)$  below, and possible targets for the level  $L(i+1)$  above.

Let us now consider, for any  $A \in L(i) \subset K^3$ , the component  $\mathcal{M}(A) \subset \mathcal{M}(3)$ , which is given for  $A \neq I$  by all local homogeneous moduli of 3-geometries with characteristic isometry  $A$ , and for  $A = I$  by  $\mathcal{M}(I) = \emptyset$ . With  $\dim \emptyset := -1$ , and components  $\text{Geom}(A) \subset \text{Geom}(3)$  of characteristic isometry of type  $A \in K^3$ , the relation

$$J(A) = \dim \text{Geom}(A) = 1 + \dim \mathcal{M}(A) \quad (4.9)$$

is satisfied for all  $A \in K^3$ . By Eq. (4.9) the Morse-like potential  $J$  on  $K^3$ , which was constructed purely topologically from  $\kappa^3$ , provides also a potential for the isometry components of  $\mathcal{M}(3)$  in terms of their dimension.

The Morse-like isometry potential for components  $\mathcal{M}(A) \subset \mathcal{M}(3)$  of the moduli space might be used to determine an evolution of a 3-geometry's isometry towards the minimum, corresponding to the flat geometry  $I^3$ . However, a 3-geometry in the interior of its characteristic isometry component  $\text{Geom}(A)$  with  $J(A) = i > 0$ , contained in some nonminimal potential level  $L(i)$ , might be considered as *metastabilized* against a transition: Since all isometry components of lower potential level can be reached only on the boundary of the isometry component  $\text{Geom}(A)$ , for that given geometry, a transition to lower level is inhibited by the geometry's distance to that boundary.

## 5. Discussion

With the flat center of projection  $I^3$  excluded, the moduli space  $\mathcal{M}(3)$ , of local homogenous Riemannian 3-geometries  $\text{Geom}(3)$  modulo homogeneous conformal transformations  $\text{Conf}(3)$ , has been constructed as an algebraic variety. The explicit parametrization of  $\mathcal{M}(3)$  by  $\hat{R}$ ,  $\hat{S}$ ,  $\hat{Y}$ , and of  $\text{Conf}(3)$  by  $N$ , yields invariant measures on both, and on  $\text{Geom}(3)$ , just given by the scalar invariants of geometry. Although the moduli space  $\mathcal{M}(3)$  is not locally Euclidean, it is a Hausdorff space. Nevertheless, Eq. (4.9) shows that, its topology is also compatible with the non-Hausdorffian topology  $\kappa^3$  of the space  $K^3$  of all Bianchi-Lie algebras, which characterize the moduli already up to differences in their anisotropic scales. Independently from regularity requirements in the construction of  $\mathcal{M}(3)$ , Eq. (4.9) and its possible interpretation related to an isometry potential, show that the definition (1.9) is, at least for  $d = 3$ , the right one.

With the (perhaps a little artificial) notion of *characteristic isometry*, assigning the Bianchi Lie isometry  $IX = so(3)$  to the KS spaces, the problem of occasionally missing simply transitive isometries could be circumvented for the considered dimension  $d = 3$ . For  $d > 3$  this problem might be much worse. There, also the parametrization of  $\text{Geom}(d)$  by scalar invariants is essentially more difficult, due to the additional presence of the Weyl tensor.

The analogous construction for local homogeneous Lorentzian 3-geometries is complicated by the null cone structure, as additional datum of the geometry. In this case, partial results were obtained in [13]. The non-uniqueness of the flat Lorentzian 3-geometry implies further singularities, which have to be taken into account in a moduli construction.

For canonical quantization, in the local homogeneous case considered here, the factorization of non-flat 3-geometries, into conformal modes  $\text{Conf}(3)$  and moduli  $\mathcal{M}(3)$ , provides both, a possible regularization of an Euclidean path integral over 3-geometries, and the foundation for a conformal mode quantization with fixed moduli. The latter technique is extensively used in minispace constructions of multidimensional geometries (see e.g. [3, 4, 17, 18] and references therein).

Although global properties of the homogeneous 3-geometries have not been considered here, it should be clear that the global properties are partially dependent on

the local ones: Any global homogeneous 3-geometry of a given Thurston type, admits only very specific local geometries (cf. [19]), corresponding to characteristic points in the moduli space  $\mathcal{M}(3)$ .

**Acknowledgement:** The author thanks the DAAD for financial support of this work, and the IPM for hospitality and for supporting the participation at the Pacific Conference of Gravitation and Cosmology. Comments of H.-J. Schmidt are gratefully acknowledged. M. Mohazzab deserves special thanks for helpful discussions and figure plotting.

## References

- [1] M. Visser, Phys. Rev. D **42**, 1964-1972 (1990).
- [2] H.-J. Schmidt, p. 405 in: Diff. Geom. and Appl., ed.: D. Krupka, Brno 1989.
- [3] M. Rainer, Gravitation & Cosmology **1**, 121-130 (1995).
- [4] M. Rainer, Int. J. Mod. Phys. D **4**, 397-416 (1995).
- [5] M. Rainer, J. Math. Phys. **35**, 646-655 (1994).
- [6] A. E. Fischer, Gen. Rel. Grav. **15**, 1191 (1983); J. Math. Phys. **27**, 718 (1986).
- [7] S. T. Swift, J. Math. Phys. **33**, 3723 (1992).
- [8] J. D. Brown and E. A. Martinez, Phys. Rev. D **42**, 1931-1943 (1990).
- [9] L. D. Landau and E. M. Lifschitz, *Lehrbuch der theoretischen Physik II*, 12. Aufl., pp. 457 ff. Akademie-Verlag, Berlin (1992).
- [10] D. Kramer, H. Stephani, M. MacCallum and E. Herlt, *Exact solutions of Einstein's field equations*, Sec. 8.6, p. 106. Wissenschaften, Berlin (1980).
- [11] J. Milnor, Adv. in Math. **21**, 293-329 (1976).
- [12] K. Nomizu, Osaka J. Math. **16**, 143-150 (1979).
- [13] M. Mohazzab, M. Rainer, and H.-J. Schmidt, *Deformations between Bianchi geometries in classical and quantum cosmology*, Preprint UniP-Math-95/04 and IPM-95-91 (1995).
- [14] M. Rainer and H.-J. Schmidt, Gen. Rel. Grav. **27**, 1265-1294 (1995).
- [15] M. Rainer, *The topology of the space of real Lie algebras up to dimension 4 with applications to homogeneous cosmological models*, Dissertation, Univ. Potsdam (June 1994), publ. in: Edition Wissenschaft, Reihe: Mathematik, Bd. 3, Marburg (1995).
- [16] M. Rainer, alg-geom/9508006, to be publ. in: Algebras, Groups and Geometries (1996).
- [17] V. D. Ivashchuk, V. N. Melnikov and A. I. Zhuk, Nuovo Cim. B **104**, 575 (1989).
- [18] V.D. Ivashchuk and V.N. Melnikov, Gravitation & Cosmology **1**, 121-130 (1995).
- [19] M. Lachize-Rey and J.-P. Luminet, Physics Reports **254**, 135-214 (1995).

Fig. 1: Riemannian Bianchi geometries II, IV, V,  $\text{VI}_h(w=0)$ ,  $\text{VI}_h$  ( $\sqrt{h} = 0, \frac{1}{5}, \frac{1}{4}, \frac{1}{3}, \frac{5}{8}, 1, 2$ ),  $\text{VII}_h$  ( $\sqrt{h} = 0, \frac{1}{7}, \frac{1}{5}, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, 1$ ); w.r.t. the common origin, the axes of the 3 planar diagrams, are:  $\hat{R}/\sqrt{3}$  to the right,  $\sqrt{6}\hat{S}$  up, and  $2 \tanh \hat{Y}$  both, left and down.

Fig. 2: Riemannian Bianchi geometries II, V, VI<sub>0</sub>, VI<sub>1</sub>, VII<sub>0</sub>, VIII( $t, w$ ) ( $t = -5, -1, 0, 1, 5$ ), IX( $t, w$ ) ( $t = 0, \frac{1}{2}, 1, 2, 5$ ); w.r.t. the common origin, the axes of the 3 planar diagrams, are:  $\hat{R}/\sqrt{3}$  to the right,  $\sqrt{6}\hat{S}$  up, and  $2 \tanh \hat{Y}$  both, left and down.

Fig. 3: The topological space  $K^3$  (right and left images have to be identified for the algebras IV and V; the locally maximal algebras IV,  $\text{VI}_h$  and  $\text{VII}_h$ ,  $0 \leq h < \infty$ , form a 1-parameter set of sources of arrows) as isometry potential.



This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9607067v1>

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9607067v1>

This figure "fig1-2.png" is available in "png" format from:

<http://arXiv.org/ps/gr-qc/9607067v1>